

# The Quantum Big Bang in Global Time Theory

D.E. Burlankov\*

Physics Department, University of Nizhny Novgorod, Russia.

The *Global Time Theory* (GTT) is the further development of the General Relativity (GR). GTT significantly differs from GR in the general physical concepts, but retains 90% of the mathematical structure and main results. The dynamics equations are derived from Lagrangian, and the Hamiltonian of gravitation is nonzero. The quantum theory of gravitation can be built on the basis of the Schroedinger equation, as for other fields. The quantum model of the Big Bang is demonstrated.

PACS numbers: 04, 04.60.-m, 98.80.Hw

## I. GLOBAL TIME THEORY

The Global Time Theory (GTT) (see [1]) is the further development of General Relativity (GR). GTT significantly differs from GR in base physical postulates, but retains 90% of the mathematical structure and main results. Meanwhile, description of the cosmos dynamics in GTT leads to significant modifications.

In GTT time is absolute. It flows equally – always and everywhere – and is itself the measure of equality. The development of the *Entire Universe* occurs in this global time.

The space has three dimension, is Riemanian, and its metric tensor ( $\gamma_{ij}$ ) can depend on space coordinates and time. Points of space are linked with it absolutely. The frame of reference in which coordinates of space points do not change is called the *global inertial system*.

The inertial system permits arbitrary tree-dimensional transformations of coordinates  $\tilde{x}^i(x^j)$  (that do not depend on time).

The coordinate transformations that are *time-dependent* lead to *global non-inertial system* of observer. Meanwhile the time remain global. In the non-inertial system the *vector field of absolute velocities*  $V^i$  arises, although it vanish in the inertial system.

Thus there are nine components of fields, described the space dynamics: six components of three-metric  $\gamma_{ij}$  and three components of absolute velocities  $V^i$ .

The space dynamics describes in global time but usually in *non-inertial system* which coordinates  $x^i$  are linked with inertial coordinates  $\bar{x}^i$  by transformations, depending on the global time  $t$ . As sequence of this transformations the time derivative in inertial system is expressed through the derivatives in the non-inertial system by the it covariant derivative over the time. The structure of covariant derivatives over the time of tensors acquires additions in the form of Lie-variation, that are generated by transformation of coordinates  $\delta x^i = -V^i dt$  – for return to the inertial system.

For a tensor of an arbitrary range

$$D_t Q_{jk}^i = \frac{\partial}{\partial t} Q_{jk}^i - V_{;s}^i Q_{jk}^s + V_{;j}^s Q_{sk}^i + V_{;k}^s Q_{js}^i + V^s Q_{jk;s}^i. \quad (1)$$

For scalar there is

$$D_t F = \frac{\partial F}{\partial t} + \frac{\partial F}{\partial x^i} \frac{\partial x^i}{\partial t} = \frac{\partial F}{\partial t} + V^i \frac{\partial F}{\partial x^i}, \quad (2)$$

which determines the covariant derivative over the time of scalar field (action, eikonal) in an arbitrary frame with global time.

Especially important for the theory is the covariant derivative over the time of the metric tensor:

$$D_t \gamma_{ij} = \frac{\partial \gamma_{ij}}{\partial t} + V_{i;j} + V_{j;i}. \quad (3)$$

### A. Action and dynamical equations

In GR, space play a rather passive role. In GTT the three-dimensional space is the dynamic field, relative to which there exists an absolute motion, or, on the contrary, there exists a field of space velocities in a given system of coordinates. The equations of motion are derived from the variational principle. Lagrangian is presented as the difference of the kinetic and potential energy. Introducing the *tensor of space deformation velocity*

$$\mu_{ij} = \frac{1}{2c} D_t \gamma_{ij} = \frac{1}{2c} (\dot{\gamma}_{ij} + V_{i;j} + V_{j;i}), \quad (4)$$

we can represent this action as preconcerted with the Hilbert action in GR:

$$S = \frac{c^4}{16\pi k} \int (\mu_j^i \mu_i^j - (\mu_j^j)^2 + R) \sqrt{\gamma} d_3 x dt + S_m, \quad (5)$$

where  $S_m$  is the action of enclosed matter, which adds to dynamic equations the energy-momenta tensor components. The absolute velocities  $V^i$  are represented only in kinetic energy.

By introducing momenta

$$\pi_j^i = \frac{\sqrt{\gamma}}{2} (\mu_j^i - \delta_j^i \mu_s^s),$$

\*Electronic address: bur@phys.unn.runnet.ru

and varying the action by six components of spacial metrics, we obtain the six equations of dynamics:

$$\dot{\pi}_j^i = b_j^i + \sqrt{\gamma} G_j^i + \sqrt{\gamma} (T_j^i - V^i T_j^0), \quad (6)$$

where  $b_j^i$  we call *the self tensor current*

$$b_j^i = -\delta_j^i \frac{1}{\sqrt{\gamma}} (2 \pi_l^k \pi_k^l - \pi_k^k \pi_l^l) - \partial_s (V^s \pi_j^i) + V^i_{,s} \pi_j^s - V^s_{,j} \pi_s^i, \quad (7)$$

$G_j^i$  is the Einstein's tensor of three-dimensional space, and  $T_\beta^\alpha$  – components of tensor energy-momenta of enclosed matter, which determine *the exterior tensor current*.

In three-dimensional space, the Riemann-Kristoffel tensor is algebraically expressed through the Einstein's tensor ( $\epsilon_{ijk}$  – absolute antisymmetric tensor):

$$R_{kl}^{ij} = -\epsilon^{ijs} \epsilon_{klm} G_s^m,$$

and therefore the absence of currents leads to a flat space.

The variation (only the kinetic part of action) by three components of the field of absolute velocities gives three equations of links:

$$\nabla_i \pi_j^i = \sqrt{\gamma} \frac{8\pi k}{c^4} T_j^0. \quad (8)$$

The Hamiltonian :

$$H = \int \pi^{ij} \dot{\gamma}_{ij} d_3x - L =$$

$$\int \left( \frac{c^4}{16\pi k} \left( \frac{\pi_j^i \pi_i^j - (\pi_i^i)^2 / 2}{\sqrt{\gamma}} - R \sqrt{\gamma} \right) - 2 \pi_j^i V_{;i}^j \right) d_3x. \quad (9)$$

Its unique feature is the non fixed sign, as a result of which such phenomena as Friedman cosmological expansion are possible.

### B. The proper time of the moving body

Similar to GR, the GTT encompasses special relativity. On level with global time, in which the growth of Entire Universe occur, for the moving observer there are his local system and *local time*. All local phenomena in the moving system develop in this local time. This can be expressed through the square of absolute velocity

$$d\tau = dt \sqrt{1 - \gamma_{ij} (\dot{x}^i - V^i)(\dot{x}^j - V^j)}. \quad (10)$$

This expression can be represented in four-dimensions by joining the time and space into unified four-dimensional manifold with metrics

$$g_{00} = 1 - \gamma_{ij} V^i V^j; \quad g_{0i} = \gamma_{ij} V^j; \quad g_{ij} = -\gamma_{ij}. \quad (11)$$

The reverse metric tensor of this four-dimensional manifold is

$$g^{00} = 1; \quad g^{0i} = V^i; \quad g^{ij} = V^i V^j - \gamma^{ij}.$$

The first equation carries great significance

$$g^{00} = 1. \quad (12)$$

This is *the main structural relationship* in GTT, analogous to the Minkowski metric, which is the main structural relationship in special relativity.

### C. General Relativity

If there is the four-dimensional metric  $g_{\alpha\beta}$  in arbitrary four coordinates  $x^\alpha$ ,  $\alpha = 0..3$ , the variable  $\tau$  must be determined for reduction to global time, in order for the main structure relation (12) to hold true. We must transfer the metrics component  $g^{00}$  by rule of tensor transformation:

$$\bar{g}^{00} = g^{\alpha\beta} \frac{\partial \tau}{\partial x^\alpha} \frac{\partial \tau}{\partial x^\beta} = 1. \quad (13)$$

This differential equation turns out to be Hamilton-Jacoby differential equation for free-falling bodies (laboratories), the common time for which is  $\tau$ , which is the global time.

Thus *the equivalence principle* is realized, but in contrast to GR, the time of the inertial system exists not only for a local laboratory, but for a great many laboratories in all of space.

GTT does not mathematically coincide with GR only in one equation: since the main structure relation (12)  $g^{00} = 1$  not permit to vary this component, the tenth Einstein's equation, determined by this component variation in GTT is absent. This tents equation of GR in GTT represents as

$$H = 0, \quad (14)$$

where  $H$  is the common density of the Hamiltonian of space and enclosed matter.

Thus the GR solutions is the subset of GTT solutions with particular value of energy equal to zero.

## II. BIG BANG IN GTT

So as the Hamiltonian in GTT is not equal zero, the effective quantum theory of gravitation can be built on the basis of the Schroedinger equation, as for other fields.

### A. Classical solutions

Now we study the compact cosmological model of Friedman type with space as a three-dimensional sphere with variable radius  $r$ , depended on the time  $t$ . This Universe is filled by ultra-relativistic matter with the state equation  $\varepsilon = 3p$ .

The Lagrangian of isoentrophic gas is expressed by integral over space of pressure, determined as function of the chemical potential [2]  $\mu$  as function (in homogeneous case) of the time derivative of *inner action*:  $\mu = \dot{\sigma}$ :

$$L_m = 2\pi^2 \int p(\mu) r^3. \quad (15)$$

The action variation by  $\sigma$  yields the matter conservation law

$$\frac{d}{dt}(\rho r^3) = 0; \quad \rho = \frac{dp}{d\mu}. \quad (16)$$

For ultrarelativistic matter the pressure is proportional  $\mu^4$ , what together with Lagrangian of space yields the full Lagrangian:

$$L = -\frac{3\pi c^2}{4k} r \dot{r}^2 + \frac{3\pi c^4 r}{4k} + \frac{\dot{\sigma}^4}{4a^6} r^3, \quad (17)$$

where constant  $a$  has dimension of length.

Further we will work in almost Planckian system of units where light velocity  $c = 1$ ,  $2k/(3\pi) = 1$  and  $\hbar = 1$ . All physical values are non-dimensional and the dimensional energy  $E$  express through non-dimensional  $e$  by expression

$$E = e c^2 \sqrt{\frac{3\pi c \hbar}{2k}}.$$

The non-dimensional Lagrangian is

$$L = \frac{r(1 - \dot{r}^2)}{2} + \frac{\dot{\sigma}^4}{4a^6} r^3.$$

The momentum conjugated to  $\sigma$  is constant

$$p_r = -r \dot{r}; \quad p_\sigma = \frac{\dot{\sigma}^3 r^3}{a^6}.$$

By expression the velocities through momenta

$$\dot{r} = -\frac{p_r}{r}; \quad \dot{\sigma} = \frac{1}{r} \left( \frac{p_\sigma}{\rho_0} \right)^{1/3},$$

we can express the Hamiltonian

$$H = -\dot{r} p_r + \dot{\sigma} p_\sigma - L = -\frac{p_r^2 + r^2}{2r} + \frac{q^2}{2r}, \quad (18)$$

where  $q = \sqrt{3/2} a p_\sigma^{(2/3)}$  and determines the conserved quantity of the ultrarelativistic matter.

The classical equation of motion can be derived from the energy conservation law:

$$\dot{r} = \frac{1}{r} \sqrt{q^2 - 2Er - r^2}. \quad (19)$$

This equation describes the radius oscillations between the maximal and minimal values, which are determined by roots of the subroot expression.

$$r_{max} = \sqrt{e^2 + q^2} - e. \quad (20)$$

The second root is negative.

If  $q^2 \neq 0$ , the energy can be as negative as positive. At  $q^2 = 0$  – clear gravitational dynamics without matter – the energy can attain only negative values,

## B. Quantum model

The wave function is function of the radius  $r$ . By  $u'$  we denote the derivative of the wave function  $u(r)$  over  $r$ . By symmetrization the product  $p^2/r$  we get the cosmological wave equations:

$$-\left(\frac{u'}{r}\right)' + \left(-r + \frac{q^2}{r}\right) u = 2Eu.$$

$$u'' - \frac{u'}{r} + (-r^2 + q^2) u = 2Eu : \quad (21)$$

This equation has a regular critical point  $r = 0$  and irregular  $r = \infty$ , at that vicinity the wave function behave oneself as the ones of oscillator:

$$u(r) \sim A e^{-r^2/2} + B e^{r^2/2}.$$

At the some values of  $E$  the coefficient  $B$  vanishes – there are normalizable solutions of the quantum equation. The function is equal zero at  $r = 0$  and infinity, and consequently it can have  $n$  extremums.

At the vicinity of zero radius all solutions behave oneself as  $r^2$ . This means that probability density at  $r = 0$  in any state is zero.

The equation (21) have two parameters:  $q$  and  $e$ .

The self energy values for small  $n$  at  $q^2 = 0, 1, 10$  are represented in next table

| n   | $q^2 = 0$ | $q^2 = 1$ | $q^2 = 10$ |
|-----|-----------|-----------|------------|
| 1   | -1.3133   | -1.0202   | 2.6765     |
| 2   | -1.9243   | -1.7122   | 0.564      |
| 3   | -2.3863   | -2.2107   | -0.441     |
| 4   | -2.773    | -2.6193   | -1.1208    |
| ... | ...       | ...       | ...        |
| 8   | -3.9599   | -3.8487   | -2.8153    |

The eigenvalues of energy can be equal to zero (as in GR), but only at the  $q^2 = 4n$ . At this value of  $q$  there is solely wave function and only with  $n$  extremums.

This model demonstrate the difference between quantum and classical solutions. The solutions of the classical

equation (19) describes the radius oscillations between the positive maximal and negative minimal radiuses. The point  $r = 0$  in classical solution is usual point. In quantum solutions the point  $r = 0$  is critical point and the value of the wave function at this point is zero, so the quantum oscillations of radius take place between zero and maximal classical radius.

### III. CONCLUSION

The main philosophical difference of GTT from GR is in its rejection of the *principle of general covari-*

*ance*. This difference is not in its mathematical treatment (upon coordinate transformation all equations must be transformed correctly), but in its physical vulgarization, i.e the confirmation of the principal *physical* equality of all nonsingular coordinate systems, and as a result – the confirmation of principal nonpossibility of the introduction of global time. GTT revives the global time, which simplifies and enriches the theory.

The main *physical* difference of GTT from GT is in nonzero energy density. In contrast to GR, in GTT the quantum theory of gravitation is built on the basis of the Schroedinger equation, as for other fields.

Author thanks Ksenia P. Brazhnik for help in translation.

- 
- [1] D.E. Burlankov. Physics-Uspekhi, **174** N 7 (2004)
  - [2] B.F. Schutz. Phys. Rev. D, **4**, 3559-3566 (1971)